

References for Component Testing for Food Allergy Diagnosis

Ando H, Movérare R, Kondo Y, et al. Utility of ovomucoid-specific IgE concentrations in predicting symptomatic egg allergy. *J Allergy Clin Immunol*. 2008;122(3):583-588.

doi: [10.1016/j.jaci.2008.06.016](https://doi.org/10.1016/j.jaci.2008.06.016)

Bartnikas LM, Sheehan WJ, Larabee KS, Petty C, Schneider LC, Phipatanakul W. Ovomucoid is not superior to egg white testing in predicting tolerance to baked egg. *J Allergy Clin Immunol Pract*. 2013;1(4):354-60. doi: [10.1016/j.jaip.2013.04.002](https://doi.org/10.1016/j.jaip.2013.04.002)

Bégin P, Vitte J, Paradis L, et al. Long-term prognostic value of component-resolved diagnosis in infants and toddlers with peanut allergy. *Pediatr Allergy Immunol*. 2014;25(5):506-8.

doi: [10.1111/pai.12216](https://doi.org/10.1111/pai.12216)

Berneer M, Bublin M, Hoffmann-Sommergruber K, et al. Allergen chip diagnosis for soy-allergic patients: Gly m 4 as a marker for severe food-allergic reactions to soy. *Int Arch Allergy Immunol*. 2013;161(3):229-33. doi: [10.1159/000345970](https://doi.org/10.1159/000345970)

Beyer K, Ellman-Grunther L, Järvinen KM, Wood RA, Hourihane J, Sampson HA. Measurement of peptide-specific IgE as an additional tool in identifying patients with clinical reactivity to peanuts. *J Allergy Clin Immunol*. 2003;112(1):202-7. doi: [10.1067/mai.2003.1621](https://doi.org/10.1067/mai.2003.1621)

Beyer K, Grishina G, Bardina L, Grishin A, Sampson HA. Identification of an 11S globulin as a major hazelnut food allergen in hazelnut-induced systemic reactions. *J Allergy Clin Immunol*. 2002;110(3):517-523. doi: [10.1067/mai.2002.127434](https://doi.org/10.1067/mai.2002.127434)

Bloom KA, Huang FR, Bencharitiwong R, Nowak-Wegrzyn A, Sampson HA. Effect of Heating on Cow's Milk and Differences in Immunoblot Reactivity to Incrementally Heated Milk Among Cow's Milk-Allergic Children. *J Allergy Clin Immunol*. 2009;123(2):S182.

doi: [10.1016/j.jaci.2008.12.688](https://doi.org/10.1016/j.jaci.2008.12.688)

Bock AS, Munoz-Furlong A, Sampson HA. Further fatalities caused by anaphylactic reactions to food, 2001-2006. *J Allergy Clin Immunol*. 2007;119(4):1016-18.

doi: [10.1016/j.jaci.2006.12.622](https://doi.org/10.1016/j.jaci.2006.12.622)

Boyano-Martínez T, García-Ara C, Pedrosa M, Díaz-Pena JM, Quirce S. Accidental allergic reactions in children allergic to cow's milk proteins. *J Allergy Clin Immunol*. 2009;123(4):883-888. doi: [10.1016/j.jaci.2008.12.1125](https://doi.org/10.1016/j.jaci.2008.12.1125)

Boyano-Martínez T, Pedrosa M, Belver T, Quirce S, García-Ara C. Peach allergy in Spanish children: tolerance to the pulp and molecular sensitization profile. *Pediatr Allergy Immunol*. 2013;24(2):168-72. doi: [10.1111/pai.12037](https://doi.org/10.1111/pai.12037)

Bublin M, Breiteneder H. Cross-reactivity of peanut allergens. *Curr Allergy Asthma Rep*. 2014;14(4):426. doi: [10.1007/s11882-014-0426-8](https://doi.org/10.1007/s11882-014-0426-8)

Caubet JC, Bencharitiwong R, Moshier E, Godbold JH, Sampson HA, Nowak-Wegrzyn A. Significance of ovomucoid- and ovalbumin-specific IgE/IgG(4) ratios in egg allergy. *J Allergy Clin Immunol Pract*. 2012;129(3):739-47. doi: [10.1016/j.jaci.2011.11.053](https://doi.org/10.1016/j.jaci.2011.11.053)

D'urbano LE, Pellegrino K, Artesani MC, et al. Performance of a component-based allergen-microarray in the diagnosis of cow's milk and hen's egg allergy. *Clin Exp Allergy*. 2010;40(10):1561-70. doi: [10.1111/j.1365-2222.2010.03568.x](https://doi.org/10.1111/j.1365-2222.2010.03568.x)

Dang TD, Tang M, Choo S, et al. Increasing the accuracy of peanut allergy diagnosis by using Ara h 2. *J Allergy Clin Immunol*. 2012;129(4):1056-63. doi: [10.1016/j.jaci.2012.01.056](https://doi.org/10.1016/j.jaci.2012.01.056)

Daul CB, Slattery M, Reese G, Lehrer SB. Identification of the major brown shrimp (*Penaeus aztecus*) allergen as the muscle protein tropomyosin. *Int Arch Allergy Immunol*. 1994;105:49-55. doi: [10.1159/000236802](https://doi.org/10.1159/000236802)

Ebisawa M, Brostedt P, Sjolander S, et al. Gly m 2S albumin is a major allergen with a high diagnostic value in soybean-allergic children. *J Allergy Clin Immunol*. 2013;132(4):976-8. doi: [10.1016/j.jaci.2013.04.028](https://doi.org/10.1016/j.jaci.2013.04.028)

Eller E, Bindslev-Jensen C. Clinical value of component-resolved diagnostics in peanut-allergic patients. *Allergy*. 2013;68(2):190-4. doi: [10.1111/all.12075](https://doi.org/10.1111/all.12075)

Fernandes J, Reshef A, Patton L, Ayuso R, Reese G, Lehrer SB. Immunoglobulin E antibody reactivity to the major shrimp allergen, tropomyosin, in unexposed Orthodox Jews. *Clin Exp Allergy*. 2003;33(7):956-61. doi: [10.1046/j.1365-2222.2003.01722.x](https://doi.org/10.1046/j.1365-2222.2003.01722.x)

Fernández-Rivas M, Bolhaar S, González-Mancebo E, et al. Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. *J Allergy Clin Immunol*. 2006;118(2):481-8. doi: [10.1016/j.jaci.2006.05.012](https://doi.org/10.1016/j.jaci.2006.05.012)

Flinterman AE, Akkerdaas JH, den Hartog Jager CF, et al. Lipid transfer protein-linked hazelnut allergy in children from a non-Mediterranean birch-endemic area. *J Allergy Clin Immunol*. 2008;121(2):423-428. doi: [10.1016/j.jaci.2007.10.009](https://doi.org/10.1016/j.jaci.2007.10.009)

Flinterman AE, Akkerdaas JH, Knulst AC, van Ree R, Pasmans SG. Hazelnut allergy: from pollen-associated mild allergy to severe anaphylactic reactions. *Curr Opin Allergy Clin Immunol*. 2008;8(3):261-5. doi: [10.1097/ACI.0b013e3282ffb145](https://doi.org/10.1097/ACI.0b013e3282ffb145)

Flinterman AE, Knol EF, Lencer DA, et al. Peanut epitopes for IgE and IgG4 in peanut-sensitized children in relation to severity of peanut allergy. *J Allergy Clin Immunol*. 2008;121(3):737-43. doi: [10.1016/j.jaci.2007](https://doi.org/10.1016/j.jaci.2007)

Gámez C, Sánchez-García S, Ibáñez MD, et al. Tropomyosin IgE-positive results are a good predictor of shrimp allergy. *Allergy*. 2011;66(10):1375-83. doi: [10.1111/j.1398-9995.2011.02663.x](https://doi.org/10.1111/j.1398-9995.2011.02663.x)

Gupta RS, Springston EE, Warriar MR, et al. The Prevalence, Severity, and Distribution of Childhood Food Allergy in the United States. *Peds*. 2011;128(1):e9-17.

doi: [10.1542/peds.2011-0204](https://doi.org/10.1542/peds.2011-0204)

Haneda Y, Kando N, Yasui M, et al. Ovomucoid IgE is a better marker than egg white-specific IgE to diagnose boiled egg allergy. *J Allergy Clin Immunol*. 2012;129(6):1681-82.

doi: [10.1016/j.jaci.2012.03.041](https://doi.org/10.1016/j.jaci.2012.03.041)

Ito K, Futamura M, Movérare R, et al. The usefulness of casein-specific IgE and IgG4 antibodies in cow's milk allergic children. *Clin Mol Allergy*. 2012;10(1):1. **doi:** [10.1186/1476-7961-10-1](https://doi.org/10.1186/1476-7961-10-1)

Ito K, Sjölander S, Sato S, et al. IgE to Gly m 5 and Gly m 6 is associated with severe allergic reactions to soybean in Japanese children. *J Allergy Clin Immunol*. 2011;128(3):673-5.

doi: [10.1016/j.jaci.2011.04.025](https://doi.org/10.1016/j.jaci.2011.04.025)

Järvinen KM, Beyer K, Vila L, Bardina L, Mishoe M, Sampson HA. Specificity of IgE antibodies to sequential epitopes of hen's egg ovomucoid as a marker for persistence of egg allergy. *Allergy*. 2007;62(7):758-765. **doi:** [10.1111/j.1398-9995.2007.01332.x](https://doi.org/10.1111/j.1398-9995.2007.01332.x)

Järvinen KM, Beyer K, Vila L, Chatchatee P, Busse PJ, Sampson HA. B-cell epitopes as a screening instrument for persistent cow's milk allergy. *J Allergy Clin Immunol*. 2002;110(2):293-7.

doi: [10.1067/mai.2002.126080](https://doi.org/10.1067/mai.2002.126080)

Klemans RJ, Knol EF, Michelsen-Huisman A, et al. Components in soy allergy diagnostics: Gly m 2S albumin has the best diagnostic value in adults. *Allergy*. 2013;68(11):1396-402.

doi: [10.1111/all.12259](https://doi.org/10.1111/all.12259)

Koppelman SJ, Wensing M, Ertmann M, Knulst AC, Knol EF. Relevance of Ara h 1, Ara h 2, and Ara h 3 in peanut-allergic patients, as determined by immunoglobulin E Western blotting, basophil-histamine release and intracutaneous testing: Ara h 2 is the most important peanut allergen. *Clin Exp Allergy*. 2004;34(4):583-90. **doi:** [10.1111/j.1365-2222.2004.1923.x](https://doi.org/10.1111/j.1365-2222.2004.1923.x)

Krause S, Reese G, Randow S, et al. Lipid transfer protein (Ara h 9) as a new peanut allergen relevant for a Mediterranean allergic population. *J Allergy Clin Immunol*. 2009;124(4):771-8.e5.

doi: [10.1016/j.jaci.2009.06.008](https://doi.org/10.1016/j.jaci.2009.06.008)

Lemon-Mule H, Sampson HA, Sicherer SH, et al. Immunologic changes in children with egg allergy ingesting extensively heated egg. *J Allergy Clin Immunol*. 2008;122(5):977-83 e1.

doi: [10.1016/j.jaci.2008.09.007](https://doi.org/10.1016/j.jaci.2008.09.007)

Lemon-Mulé H, Sampson HA, Sicherer SH, Shreffler WG, Noone S, Nowak-Wegrzyn A. Immunologic changes in children with egg allergy ingesting extensively heated egg. *J Allergy Clin Immunol*. 2008;122(5):977-83.e1. **doi:** [10.1016/j.jaci.2008.09.007](https://doi.org/10.1016/j.jaci.2008.09.007)

Leung PS, Chow WK, Duffey S, Kwan HS, Gershwin ME, Chu KH. IgE reactivity against a cross-reactive allergen in crustacean and mollusca: Evidence for tropomyosin as the common allergen. *J Allergy Clin Immunol*. 1996;98(5 part 1):954-961.

doi: [10.1016/S0091-6749\(96\)80012-1](https://doi.org/10.1016/S0091-6749(96)80012-1)

Lieberman JA, Huang FR, Sampson HA, Nowak-Wegrzyn. Outcomes of 100 consecutive open, baked-egg oral food challenges in the allergy office. *J Allergy Clin Immunol*. 2012;129(6):1682-4.e2. doi: [10.1016/j.jaci.2012.04.007](https://doi.org/10.1016/j.jaci.2012.04.007)

Masthoff LJ, Mattsoon L, Zuidmeer-Jongejan L, et al. Sensitization to Cor a 9 and Cor a 14 is highly specific for hazelnut allergy with objective symptoms in Dutch children and adults. *J Allergy Clin Immunol*. 2013;132(2):393-9. doi: [10.1016/j.jaci.2013.02.024](https://doi.org/10.1016/j.jaci.2013.02.024)

Matsuo H, Dahlstrom J, Tanaka A, et al. Sensitivity and specificity of recombinant omega-5 gliadin-specific IgE measurement for the diagnosis of wheat-dependent exercise-induced anaphylaxis. *Allergy*. 2008;63(2):233-236. doi: [10.1111/j.1398-9995.2007.01504.x](https://doi.org/10.1111/j.1398-9995.2007.01504.x)

Mittag D, Vieths S, Vogel L, et al. Soybean allergy in patients allergic to birch pollen: clinical investigation and molecular characterization of allergens. *J Allergy Clin Immunol*. 2004;113(1):148-54. doi: [10.1016/j.jaci.2003.09.030](https://doi.org/10.1016/j.jaci.2003.09.030)

Nicolaou N, Poorafshar M, Murray C, et al. Allergy or tolerance in children sensitized to peanut: prevalence and differentiation using component-resolved diagnostics. *J Allergy Clin Immunol*. 2010;125(1):191-7.e1-13. doi: [10.1016/j.jaci.2009.10.008](https://doi.org/10.1016/j.jaci.2009.10.008)

Nowak-Wegrzyn A, Bloom KA, Sicherer SH, et al. Tolerance to extensively heated milk in children with cow's milk allergy. *J Allergy Clin Immunol*. 2008;122(2):342-7. doi: [10.1016/j.jaci.2008.05.043](https://doi.org/10.1016/j.jaci.2008.05.043)

Palosuo K, Varjonen E, Kekki OM, et al. Wheat omega-5 gliadin is a major allergen in children with immediate allergy to ingested wheat. *J Allergy Clin Immunol*. 2001;108(4):634-8. doi: [10.1067/mai.2001.118602](https://doi.org/10.1067/mai.2001.118602)

Pascal M, Grishina G, Yang AC, et al. Molecular Diagnosis of Shrimp Allergy: Efficiency of Several Allergens to Predict Clinical Reactivity. *J Allergy Clin Immunol Pract*. 2015;3(4):521-9.e10. doi: [10.1016/j.jaip.2015.02.001](https://doi.org/10.1016/j.jaip.2015.02.001)

Pastorello EA, Vieths S, Pravettoni V, et al. Identification of hazelnut major allergens in sensitive patients with positive double-blind, placebo-controlled food challenge results. *J Allergy Clin Immunol*. 2002;109(3):563-570 doi: [10.1067/mai.2002.121946](https://doi.org/10.1067/mai.2002.121946)

Peterson A, Kull S, Rennert S, et al. Peanut defensins: Novel allergens isolated from lipophilic peanut extract. *J Allergy Clin Immunol*. 2015;136(5):1295-301.e1-5. doi: [10.1016/j.jaci.2015.04.010](https://doi.org/10.1016/j.jaci.2015.04.010)

Reese G, Ayuso R, Lehrer SB. Tropomyosin: an invertebrate pan-allergen. *Int Arch Allergy Immunol.* 1999;119(4):247-58. doi: [10.1159/000024201](https://doi.org/10.1159/000024201)

Sampson, HA. Utility of food-specific IgE concentrations in predicting symptomatic food allergy. *J Allergy Clin Immunol.* 2001;107(5):891-896. doi: [10.1067/mai.2001.114708](https://doi.org/10.1067/mai.2001.114708)

Sampson HA, Ho DG. Relationship between food-specific IgE concentrations and the risk of positive food challenges in children and adolescents. *J Allergy Clin Immunol.* 1997;100(4):444-451. <http://www.ncbi.nlm.nih.gov/pubmed/9338535>

Sicherer SH, Sampson HA. Food allergy. *J Allergy Clin Immunol.* 2010;125(2 suppl 2):S116-25. doi: [10.1016/j.jaci.2009.08.028](https://doi.org/10.1016/j.jaci.2009.08.028)

Sicherer SH, Wood RA. Advances in Diagnosing Peanut Allergy. *J Allergy Clin Immunology Pract.* 2013;1(1):1-13. doi: [10.1016/j.jaip.2012.10.004](https://doi.org/10.1016/j.jaip.2012.10.004)

Skripak JM, Matsui EC, Mudd K, Wood RA. The natural history of IgE-mediated cow's milk allergy. *J Allergy Clin Immunol.* 2007;120(5):1172-7. doi: [10.1016/j.jaci.2007.08.023](https://doi.org/10.1016/j.jaci.2007.08.023)

Soares-Weiser K, Takwoingi Y, Panesar SS, et al. The diagnosis of food allergy: a systematic review and meta-analysis. *Allergy.* 2014;69(1):76-86. doi: [10.1111/all.12333](https://doi.org/10.1111/all.12333)

Valenta R, Lidholm J, Niederberger V, Hayek B, Kraft D, Grönlund, H. The recombinant allergen-based concept of component-resolved diagnostics and immunotherapy (CRD and CRIT). *Clin Exp Allergy.* 1999;29(7):896-904. doi: [10.1046/j.1365-2222.1999.00653.x](https://doi.org/10.1046/j.1365-2222.1999.00653.x)

Vereda A, van Hage M, Ahlstedt S, et al. Peanut allergy: Clinical and immunologic differences among patients from 3 different geographic regions. *J Allergy Clin Immunol.* 2011;127(3):603-7. doi: [10.1016/j.jaci.2010.09.010](https://doi.org/10.1016/j.jaci.2010.09.010)

Wal JM. Bovine milk allergenicity. *Ann Allergy Asthma Immunol.* 2004;93(5 Suppl 3):S2-11. <http://www.ncbi.nlm.nih.gov/pubmed/15562868>

Wang J, Calatroni A, Visness CM, Sampson HA. Correlation of specific IgE to shrimp with cockroach and dust mite exposure and sensitization in an inner-city population. *J Allergy Clin Immunol.* 2011;128(4):834-7. doi: [10.1016/j.jaci.2011.07.045](https://doi.org/10.1016/j.jaci.2011.07.045)

Yang AC, Arruda LK, Santos AB, et al. Measurement of IgE antibodies to shrimp tropomyosin is superior to skin prick testing with commercial extract and measurement of IgE to shrimp for predicting clinically relevant allergic reactions after shrimp ingestion. *J Allergy Clin Immunol.* 2010;125(4):872-8. doi: [10.1016/j.jaci.2009.11.043](https://doi.org/10.1016/j.jaci.2009.11.043)